

The range of validity of cluster masses and ages derived from broad-band photometry

J. Maíz Apellániz¹

¹Instituto de Astrofísica de Andalucía-CSIC, Granada, Spain
 email: jmaiz@iaa.es

Abstract. I analyze the stochastic effects introduced by the sampling of the stellar initial mass function (SIMF) in the derivation of the individual masses and the cluster mass function (CMF) from broad-band visible-NIR unresolved photometry. The classical method of using unweighted *UBV* photometry to simultaneously establish ages and extinctions of stellar clusters is found to be unreliable for clusters older than ≈ 30 Ma, even for relatively large cluster masses. On the other hand, augmenting the filter set to include longer-wavelength filters and using weights for each filter increases the range of masses and ages that can be accurately measured with unresolved photometry. Nevertheless, a relatively large range of masses and ages is found to be dominated by SIMF sampling effects that render the observed masses useless, even when using *UBVR1JHK* photometry.

Keywords. methods: analytical — methods: numerical — methods: statistical — open clusters and associations: general — globular clusters: general — galaxies: star clusters

1. Description

This work is the third of a series in which we are analyzing the possible biases present in mass functions (Maíz Apellániz & Úbeda 2005; Maíz Apellániz 2008). I have used a combination of analytical approximations and Monte Carlo simulations to study the effect of the stochastic sampling of the stellar IMF (SIMF, assumed to be Kroupa) in the determination of masses and ages of unresolved stellar clusters from broad-band photometry. For this purpose I have used the new (3.1) version of CHORIZOS (Maíz Apellániz 2004), which incorporates an evolutionary synthesis module. For a given mass and age, I generated a minimum of 10 000 realizations of the SIMF using solar metallicity [a] Geneva isochrones for massive stars and [b] Padova isochrones with new AGB treatment for low and intermediate masses. In each case I measured the mass (and age, if appropriate) assuming that the cluster has a well-sampled SIMF. From the individual realizations I derived [a] the observed cluster mass and age distributions for clusters of a (fixed) real mass and age and [b] the observed cluster mass functions (CMFs) for a real truncated power-law CMF with a slope of $\gamma = -2.0$. Three different cases in order of complexity were considered: [a] single-filter observations of clusters with known age and extinction, [b] multi-filter observations of clusters with unknown age and known extinction, and [c] multi-filter observations of clusters with unknown age and extinction. In the first case masses were computed directly by converting filter-convolved luminosities to masses while in the second and third cases a χ^2 minimization code (CHORIZOS) was used to derive the masses and ages. The reader is referred to Maíz Apellániz (2009) for further details.




Figure 1. Results of the Monte Carlo simulations of $f_{M_o|M}(m_o|m)$ and Gaussian and Gamma fits for 10 Ma clusters observed in V with four real masses: $300 M_\odot$ (upper left), $1000 M_\odot$ (upper right), $10\,000 M_\odot$ (lower left), and $100\,000 M_\odot$ (lower right). The Poisson mass θ for this filter and age is $566 M_\odot$. For large real masses, both fits provide reasonable approximations but for small values of m , the Gamma approximation is significantly better. The vertical dotted lines show the values of the real mass m and, when visible, of θ and $m_{o,LLL}$ (the observed cluster mass obtained when one mistakes the brightest possible star in the isochrone for a cluster).

2. Case 1: Single filter, known age and extinction

- For a given cluster mass (m), the observed cluster mass (m_o) distribution can be approximated by a Gamma function better than by a Gaussian distribution (Figure 1):

$$f_{M_o|M}(m_o|m) = A m_o^{m/\theta - 1} e^{-m_o/\theta} \theta^{-m/\theta}. \quad (2.1)$$

- The Poisson mass θ is a measurement of the stochasticity and (strongly) depends on the age and the filter used to derive m_o (Table 2).

$$\overline{m_o} = m \quad ; \quad \sigma_{m_o} = \sqrt{\theta m}. \quad (2.2)$$

- The large differences in θ are a consequence of the different fraction of stars located above the median mass for the luminosity in each filter for a given isochrone (Figure 2).
- The observed CMF, $f_{M_o}(m_o)$, differs from the real (truncated power-law) CMF,




Figure 2. Isochrones for seven of the nine ages used in this work. For each isochrone, symbols of different colors are used to indicate the median mass for the luminosity in the eight filters U , B , V , R , I , J , H , and K assuming a Kroupa IMF between $0.1 M_{\odot}$ and $120 M_{\odot}$. The two dashed lines join the points in each isochrone beyond which 0.2% and 0.1%, respectively, of the remaining stars in a well-sampled Kroupa IMF are located. The two remaining isochrones are not included for the sake of clarity.

$f_M(m)$, in two aspects (Figure 3): [a] For large masses, $f_{M_o}(m_o)$ is similar to $f_M(m)$ but, as the cluster mass decreases, at one point a "hump" or overdensity appears. [b] To the left of the hump, $f_{M_o}(m_o)$ asymptotically approaches a power law with a slope larger than -1.0 , with some clusters having m_o values lower than the real cutoff in m .

• Using an analytical approximation, it can be shown that if one is willing to tolerate a systematic error in the CMF slope γ up to $\Delta\gamma$, then one can only analyze observed masses:

$$m_o > 0.5\gamma(\gamma + 1)\theta/\Delta\gamma, \quad (2.3)$$

e.g. for $\gamma = -2.0$ and $\Delta\gamma = 0.1$, $m_o > 10\theta$.

Table 1. Values of θ (in $M_{\odot}$) for different ages and filters for clusters observed with a single filter and known age and extinction. The labels below each age indicate the type of the brightest possible star. For a given filter, θ starts at a low value in the MS phase, rapidly grows in the LBV phase, decreases during the RSG and early AGB phases, and then experiences a moderate growth when the cluster becomes old. For a given age, θ increases as a function of wavelength with the only exception of the MS phase, where the effect is the opposite (but rather weak).

	1 Ma	3.16 Ma	10 Ma	31.6 Ma	100 Ma	316 Ma	1 Ga	3.16 Ga	10 Ga
Filter	MS	LBV	RSG	AGB1	AGB2	AGB3	AGB4	AGB5	AGB6
<i>U</i>	235	945	160	149	53	33	23	26	42
<i>B</i>	200	1 314	192	305	51	32	27	52	92
<i>V</i>	190	1 555	566	370	89	38	51	106	163
<i>R</i>	186	1 722	1 034	443	151	72	109	173	242
<i>I</i>	178	1 949	1 585	616	273	463	476	477	581
<i>J</i>	157	2 741	2 240	1 170	708	3 598	2 930	2 702	2 445
<i>H</i>	146	3 081	2 476	1 640	1 281	5 184	4 828	4 499	4 123
<i>K</i>	140	3 333	2 542	1 771	1 527	5 717	5 688	5 512	5 058




Figure 3. Observed (continuous lines) and real (dotted lines) CMFs for the minimum and maximum values of θ in Table 2. The left panel corresponds to 1 Ga clusters observed with *U* and the right panel to 316 Ma clusters observed with *K*. Each panel shows four cases of the lower mass cutoff for $f_M(m)$ from $1 M_{\odot}$ to $1000 M_{\odot}$. As θ increases, the hump becomes more pronounced and moves towards the right.

3. Case 2: Multiple filters, unknown age, and known extinction

I used three execution types:

1. *UBVR IJHK* photometry with single-band θ -dependent weights for each filter.
2. *UBVR IJHK* photometry with constant weights.
3. *UBV* photometry with single-band θ -dependent weights for each filter.

The observed age distributions for $10^4 M_{\odot}$ clusters are shown in Figure 4. The main results are:

- The ages calculated with *UBVR IJHK* photometry and constant weights (i.e. the standard method) are highly uncertain even for $10^4 M_{\odot}$ clusters.
- Using weights based on the stochasticity of each filter significantly narrows the observed-age distribution for $10^4 M_{\odot}$ clusters, thus allowing for the obtention of ages with relatively small uncertainties.




Figure 4. Distribution of observed ages for the nine input ages and three execution types (1: upper left, 2: upper right, 3: bottom) for clusters of $10^4 M_{\odot}$ and unknown ages. At the top of each panel the dashed line indicates the real age while the symbols and error bars provide the median and inferior and superior uncertainties (1-sigma equivalents) of each distribution.

- *UBV* and *UBVRIJHK* photometry provide relatively similar results, with the former having the advantage for old clusters and the latter for intermediate-age ones.
- The AGB1 stage (31.6 Ma) has significantly larger uncertainties than the rest.

4. Case 3: Multiple filters, unknown age and extinction

I used two execution types:

1. *UBVRIJHK* photometry with single-band θ -dependent weights for each filter.
2. *UBV* photometry with single-band θ -dependent weights for each filter.

The observed age distributions for $10^4 M_{\odot}$ and $10^5 M_{\odot}$ clusters are shown in Figure 5. The main results are:

- As opposed to the cases where extinction is known, the addition of *RIJHK* photometry provides a significant improvement in the observed ages.
- *UBV* photometry alone is not sufficient to accurately determine the ages of unresolved stellar clusters, even for massive ones, if extinction is unknown.
- For some ages (1-10 Ma, 100 Ma, 10 Ga), *UBVRIJHK* photometry provides relatively accurate observed ages for $m = 10^4 M_{\odot}$. For the rest, higher masses are needed.




Figure 5. Distribution of observed ages for the nine input ages and two execution types (1: left plots, 2: right plots) for clusters of $10^4 M_{\odot}$ (top) and $10^5 M_{\odot}$ (bottom) and unknown ages and extinctions. At the top of each panel the dashed line indicates the real age while the symbols and error bars provide the median and inferior and superior uncertainties (1-sigma equivalents) of each distribution.

5. Conclusions

- SIMF sampling effects can introduce large biases in the determination of masses and ages from unresolved photometry.
- One should use weights based on the relative stochasticity of each filter when using χ^2 -minimization techniques to calculate cluster properties from unresolved photometry.
- The addition of a broad-photometric baseline (U to K) significantly reduces uncertainties when extinction is determined from the data.
- For clusters younger than 100 Ma, a critical (post-RSG) stage exists around 30 Ma where ages are especially difficult to determine.

References

- Maíz Apellániz, J. 2004, *PASP*, 116, 859
 Maíz Apellániz, J. 2008, *ApJ*, 677, 1278
 Maíz Apellániz, J. 2009, *ApJ*, 699, 1938
 Maíz Apellániz, J. & Úbeda, L. 2005, *ApJ*, 629, 873